



Effects of graded levels of cupric citrate on growth performance, antioxidant status, serum lipid metabolites and immunity, and tissue residues of trace elements in weaned pigs

Chu Cai Peng¹, Jia You Yan², Bin Dong¹, Lin Zhu¹, Yao Yao Tian¹, and Li Min Gong^{1,*}

* Corresponding Author: Li Min Gong
Tel: +86-10-6273-3590, Fax: +86-10-6273-3700,
E-mail: gonglimin@cau.edu.cn

¹ State Key Laboratory of Animal Nutrition, China
Agricultural University, Beijing 100193, China

² Institute of Animal Nutrition, Sichuan Academy of
Animal Science, Chengdu 610066, China

Submitted Mar 2, 2016; Revised Apr 27, 2016;
Accepted Jun 28, 2016

Objective: The goal of this study was to investigate the effects of cupric citrate (CuCit) on growth performance, antioxidant indices, serum lipid metabolites, serum immune indices, and tissue residues of copper (Cu), zinc, and iron in weaned pigs.

Methods: A total of 180 weaned pigs (Duroc×Landrace×Large White) with an average body weight of 8.98±1.21 kg were randomly assigned to a corn-soybean meal control ration, or 4 similar rations with 30, 60, 120, or 240 mg/kg Cu as CuCit. All diets contained 10 mg/kg Cu as cupric sulfate from the vitamin-mineral premix. The experiment was divided into two phases: 0 to 14 d (phase 1) and 15 to 28 d (phase 2).

Results: Average daily gain (ADG; linearly, $p < 0.01$) and average daily feed intake (ADFI; linearly and quadratically, $p < 0.05$) were affected by an increase in CuCit during phase 2. Overall period, ADG ($p < 0.05$) and ADFI ($p < 0.01$) were linearly increased with increasing dietary levels of CuCit. Serum malondialdehyde concentrations ($p < 0.05$) and glutathione peroxidase activity ($p < 0.01$) linearly decreased and increased respectively with an increase in CuCit. Serum levels of Cu-Zn superoxide dismutase were linearly affected with an increase in CuCit ($p < 0.01$). Hepatic malondialdehyde levels decreased with an increase in CuCit (linearly and quadratically, $p < 0.01$). Serum total cholesterol concentrations were quadratically affected ($p < 0.05$) and decreased in pigs fed Cu as CuCit at 60 and 120 mg/kg and increased in pigs fed 240 mg/kg Cu as CuCit. Serum high-density lipoprotein concentrations were linearly affected with an increase in CuCit ($p < 0.01$). Serum IL-1 β levels were quadratically affected ($p < 0.05$) by dietary treatment. Compared with other treatments, 240 mg/kg Cu from CuCit quadratically increased hepatic ($p < 0.01$) and renal ($p < 0.05$) Cu concentrations, and quadratically decreased hepatic and renal iron concentrations ($p < 0.05$).

Conclusion: Cu administered in the form of CuCit at a dosage range of 30 to 60 mg/kg, effectively enhanced the growth performance and antioxidant status of weaned pigs.

Keywords: Cupric Citrate; Performance; Serum Metabolites; Antioxidant Status; Tissue Trace Elements; Weaned Pigs

INTRODUCTION

Cupric supplementation at pharmacological doses beyond the recommendations of National Research Council has been widely applied in the feed industry to improve pig performance [1-3]. While organic copper (Cu) is more efficient than inorganic Cu in enhancing the growth performance of pigs [4,5]. Copper sulfate (CuSO₄), the most common form of Cu used in pig industry, is soluble in both water and acid. Comparatively, cupric citrate (CuCit) is poorly soluble in water, but well soluble in acid and ammonia solutions, which facilitates better Cu absorption [6]. Therefore, the bioavailability of CuCit may be higher than that of CuSO₄. Previous studies demonstrated that dietary CuCit supplementation enhanced growth performance in

weaned pigs at much lower doses than did CuSO₄ [7,8]. Thus, if lower CuCit than CuSO₄ dietary concentrations could be used, the growth-promoting effects of Cu could be maintained, and the environmental emission of Cu could be reduced.

Cu affects a wide spectrum of biological activities such as catalyzing the formation of hydroxyl radicals, which can lead to lipid peroxidation [9], and it plays a critical role in iron and zinc homeostasis [6,10]. In addition, pharmacological levels of Cu affected lipid metabolism in swine [11,12], and Cu intake, insufficiently or excessively, reduced immunity in experimental animals [13,14]. Comparably, the positive performance of weaned pigs was observed feeding diets containing 250 mg/kg CuSO₄ [3]. Previous research found that diet supplemented with 125 mg/kg Cu as CuCit stimulated growth and improved feed intake [8]. Therefore, in the present study we assumed that the optimum levels of CuCit for weaned pigs were 120 or 240 mg/kg with 10 mg/kg Cu in vitamin-mineral premix. In the meantime, we wanted to study the effects of lower dosage (30 and 60 mg/kg) of CuCit in weaned pigs. So the objective of this study was to evaluate the effects of graded levels of CuCit on growth performance, antioxidant status, serum lipid metabolites, immunity and trace elements accumulation in tissues.

MATERIALS AND METHODS

The protocol employed in this trial was approved by the Institution of Animal Care and Use Committee of China Agricultural University. The animal experiment was conducted according to the National Institutes of Health Guidelines for the care and use of experimental animals. CuCit (98.5% CuCit and 34.4% Cu) used in this experiment was provided by Sichuan Animtech Feed Co., Ltd. (Sichuan, Chengdu, China).

Animal, diet and experimental design

At 35 d of age, a total of 180 weaned pigs (Duroc×Landrace×Large White) with an average body weight of 8.98±1.21 kg were assigned by sex and body weight to five treatments with six replicates per treatment and six pigs (three barrows and three gilts) per pen in a randomized complete block design. They were treated with gradient doses of Cu as CuCit (0, 30, 60, 120, and 240 mg/kg; Table 1). Additionally, all diets contained 10 mg/kg Cu as CuSO₄ from a vitamin-mineral premix and corn meal was replaced with CuCit in the other four treatments. Pigs were placed in concrete floor pens (1.2×2.0 m) partially slatted with steel, and each pen was equipped with a self-feeder and a nipple drinker. Pigs had *ad libitum* access to both feed and water. Diets were in a mash form and formulated to meet or exceed NRC [1] requirements in two phases: 0 to 14 d (phase 1) and 15 to 28 d (phase 2). Cu concentrations in experimental diets, analyzed and presented in Table 2, were as described by Armstrong et al [8].

Table 1. Composition and chemical composition of the basal diets (as-fed basis)^{1,2)}

Items	Phase 1	Phase 2
Ingredient (%)		
Corn meal	55.00	58.76
Soybean meal	14.32	18.27
Soybean oil	1.97	2.00
Excluded soybeans	100.00	7.05
Wheat meal	3.00	3.00
Soy protein concentrate	5.10	1.35
Whey powder	6.00	6.00
Dicalcium phosphate	1.46	1.24
Limestone	0.53	0.42
Salt	0.25	0.25
L-lysine-HCl	0.53	

Performance measurements and blood sampling

On d 0, 14, and 28 of the experiment, body weight and daily feed consumption were measured for the calculation of average daily gain (ADG), average daily feed intake (ADFI), and gain-to-feed ratio (G:F). On d 28, six pigs (three barrows and three gilts) from each treatment (one pig per pen) were selected based on the mean body weight of each pen. Blood samples were collected from the precaval vein after 12 h fasting and allowed to clot at room temperature of 25°C for 20 min and centrifuged at 3,000×g for 10 min at 4°C. The serum was then removed and

Table 3. Effects of cupric citrate on growth performance in weaned pigs

Item	Copper as cupric citrate (mg/kg)					SEM	p-value		
	0	30	60	120	240		ANOVA	Linear	Quadratic
Phase 1, 0 to 14 d									
Average daily gain (g)	307	359	380	383	400	22	0.070	0.250	0.888
Average daily feed intake (g)	515	609	676	668	702	36	0.010	0.089	0.880
Gain-to-feed ratio	0.595	0.590	0.563	0.574	0.568	0.010	0.118	0.223	0.953
Phase 2, 15 to 28 d									
Average daily gain (g)	452	509	511	468	438	15	0.005	0.002	0.063
Average daily feed intake (g)	832	936	928	863	781	30	0.008	0.007	0.031
Gain-to-feed ratio	0.544	0.545	0.550	0.545	0.561	0.011	0.772	0.988	0.380
Overall period, 0 to 28 d									
Average daily gain (g)	380	434	446	426	419	16	0.084	0.023	0.323
Average daily feed intake (g)	673	773	802	767	742	26	0.029	0.009	0.161
Gain-to-feed ratio	0.563	0.563	0.555	0.556	0.564	0.009	0.874	0.845	0.346

SEM, standard error of the mean; ANOVA, analysis of variance.

Data are the means of six replicates of six pigs (three barrows and three gilts) per pen.

are shown in Table 6. Serum IL-1 β levels were quadratically affected ($p < 0.05$) by dietary treatment. There were no significant differences in serum IgM, IgA, IgG, or IFN- γ levels among all treatments.

Tissue trace elements

Table 7 shows the effects of CuCit on the concentrations of tissue trace elements in weaned pigs. There were no significant differences in serum Cu, Fe, or Zn levels among all treatments.

Compared with other treatments, 240 mg/kg Cu as CuCit quadratically increased hepatic ($p < 0.01$) and renal ($p < 0.05$) Cu concentrations, and quadratically decreased hepatic and renal Fe concentrations ($p < 0.05$). Hepatic and renal Zn levels were not affected by Cu levels.

DISCUSSION

According to the results of present study, ADG and ADFI were

Table 4. Effects of cupric citrate on antioxidant status in weaned pigs

Items	Copper as cupric citrate (mg/kg)					SEM	p-value		
	0	30	60	120	240		ANOVA	Linear	Quadratic
Serum									
Malondialdehyde (nmol/mL)	9.50	7.35	7.24	7.94	8.14	0.47	0.021	0.007	0.232
Copper-zinc superoxide dismutase (U/mL)	48.48	50.49	50.89	48.78	48.20	0.41	<0.001	<0.001	0.112
Glutathione peroxidase (U/mL)	186.54	207.72	215.74	205.41	199.12	5.55	0.017	0.003	0.133
Ceruloplasmin (U/mL)	45.78	48.43	48.53	47.91	51.07	2.18	0.570	0.629	0.524
Liver									
Malondialdehyde (nmol/mL)	6.10	3.40	3.46	4.51	6.10	0.40	<0.001	<0.001	0.001
Copper-zinc superoxide dismutase (U/mL)	30.57	28.13	27.91	29.28	29.13	0.92	0.292	0.058	0.682
Glutathione peroxidase (U/mL)	405.74	333.26	337.01	346.58	375.52	28.43	0.357	0.154	0.242
Ceruloplasmin (U/mL)	21.75	19.78	19.04	19.43	22.38	1.59	0.312	0.184	0.090

SEM, standard error of the mean; ANOVA, analysis of variance.

Data are the means of six replicates of one pig per pen.

Table 5. Effects of cupric citrate on serum concentrations of lipid metabolites in weaned pigs

Items	Copper as cupric citrate (mg/kg)					SEM	p-value		
	0	30	60	120	240		ANOVA	Linear	Quadratic
Total cholesterol (mmol/L)	1.80	1.81	1.53	1.60	1.97	0.10	0.035	0.093	0.022
Triglyceride (mmol/L)	0.45	0.52	0.46	0.47	0.33	0.05	0.141	0.550	0.057
High-density lipoprotein (mmol/L)	0.50	0.50	0.67	0.51	0.46	0.04	0.014	0.003	0.439
Low-density lipoprotein (mmol/L)	0.94	1.04	0.83	0.83	1.02	0.06	0.051	0.427	0.072
Very-low-density lipoprotein (mmol/L)	52.01	55.80	55.30	43.84	54.60	5.21	0.480	0.311	0.196

SEM, standard error of the mean; ANOVA, analysis of variance.

Data are the means of six replicates of one pig per pen.

linearly affected with increasing dietary levels of CuCit, and lower dosages (30 and 60 mg/kg) of CuCit showed a better growth performance over the experimental period from d 15 to 28 and overall period. Comparably, the dose range of CuSO_4 , which significantly improved growth performance in pigs was typically higher at 100 to 250 mg/kg [3,17,18]. Furthermore, another study predicted that peak ADG and ADFI in weaned pigs were obtained with supplemental Cu levels at 150 and 114 mg/kg, respectively, in the form of CuSO_4 [19]. However, it should be

idation, because Cu (II) ions catalyzed the formation of hydroxyl radicals, which can lead to lipid peroxidation [9]. As the component of several enzymes such as Cu/Zn-SOD [24], Cu at an appropriate dosage may decrease lipid peroxidation by increasing the activity of antioxidant enzymes [25,26]. The degree of lipid peroxidation could be evaluated by MDA [27]. Reports about the effects of pharmacological Cu in feed on tissue MDA concentrations focused on the duodenum rather than the serum and liver [16,28]. A previous study showed that duodenal MDA contents were higher when pigs were fed 225 mg/kg supplemental Cu as CuSO₄ rather than TBCC [16]. In our trial, increasing doses of CuCit led to linear and quadratic increases in serum and hepatic MDA concentrations. And Cu supplemented at doses between 30 and 120 mg/kg as CuCit led to lower serum and hepatic MDA concentrations, which was in accordance with the greater serum Cu/Zn-SOD and GSH-Px activity. Feng et al [29] found 100 mg/kg Cu increased erythrocyte Cu/Zn-SOD activity compared with that in control pigs [29]. Fry et al [28] found that pigs fed 225 mg/kg CuSO₄ had lower serum GSH-Px activity compared with that in the control treatment [28]. The lower MDA levels, higher Cu/Zn-SOD and GSH-Px activity observed in our study may be attributed to the lower Cu concentrations in tissues when pigs were fed 30 to 60 mg/kg Cu as CuCit. Thus, low dosages of Cu may decrease the formation rate of oxygen free radicals, which may increase the antioxidant levels of weaned pigs, while high Cu levels may have the opposite effect. Our research indicated the effects of dietary CuCit on hepatic oxidative stress tended to fit the hormesis curve, where hepatic MDA levels initially decreased and then increased with increasing CuCit supplementation.

There were limited reports about the effects of dietary Cu addition on serum lipid metabolites of pigs. A study suggested 100 mg/kg Cu as CuCit improved serum TC levels in weaned pigs [30], however, serum TC concentrations were highest in pigs fed the 240 mg/kg treatments. Serum HDL-C concentrations linearly increased in pigs fed CuCit at 60 mg/kg, but these treatments had no effect on TG, LDL-C, and VLDL-C levels in our trial. Li et al [31] found no Cu treatment differences were observed in plasma TC, TG, HDL-C, and LDL-C levels [31]. Therefore, the effects of Cu as CuCit on lipid metabolites in weaned pigs are inconclusive.

Cu plays an important role in maintaining the immune function of pigs, and its deficiency or excess intake could impair immune cell numbers and functions [32]. Some studies showed that interleukin-2 (IL-2) was reduced by Cu deficiency, which possibly decreased T cell proliferation [14,33]. Appropriate serum concentrations of immunoglobulins, IL-1 β , and INF- γ , which are fundamental components of the immune system, indicate the activation of an immune response [34]. In our research, serum levels of IgM, IgA, IgG, and INF- γ showed no differences among all treatments. The results were similar to recent research where 20 and 180 mg/kg Cu as CuCit had no effects on serum

IgA, IgM, and IgG levels in piglets [35]. Although serum IL-1 β levels were quadratically affected by dietary treatments in our trial, it did not indicate that CuCit had improved the serum immunity in weaned pigs. Considering the definitive mechanisms of the effects of CuCit on immunity in weaned pigs, further research is warranted.

Divalent metal transporter 1 and copper transport receptor 1 are involved in the active transport of Cu (II), Fe, and Zn (II) [36]. Thus, the absorption and utilization of Cu might be related to Fe and Zn. In addition, under certain circumstances, a mucosal block to metal absorption may occur via sequestration of Fe and Cu by ferritin and metallothionein, respectively [37,38]. In our study, pigs fed Cu at 240 mg/kg from CuCit quadratically decreased hepatic and renal Fe concentrations. There were no significant differences in serum Fe or Zn, hepatic and renal Zn levels among all treatments. Huang et al [16] indicated that hepatic Fe and Zn levels were lower and higher, respectively, in pigs fed 225 mg/kg Cu as TBCC than those in the control and those fed 225 mg/kg Cu as CuSO₄ [16]. The results were similar to those of Fry et al [28] which showed that hepatic Fe concentrations were less in TBCC pigs compared with the control group [28]. Whereas Cu lysinate (250 mg/kg) decreased Zn solubility in the digestion of broilers, supplemental CuSO₄ and TBCC did not [39]. This may indicate that diets supplemented below 120 mg/kg Cu from CuCit had no effects on serum and hepatic Fe or Zn levels when dietary Fe and Zn concentrations were supplemented above the dietary requirements in this experiment.

CONCLUSION

Cupric citrate (30 to 60 mg/kg) is an efficient copper supplement due to a relative low supplemental dosage required to maintain a high level of growth performance, antioxidant status and low tissue residues.

CONFLICT OF INTEREST

We certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

ACKNOWLEDGMENTS

This work was supported by Ministry of Agriculture of the People's Republic of China. All authors are grateful for the help in laboratory analysis by staff of Ministry of Agriculture Feed Industry Centre.

REFERENCES

1. Committee on Nutrient Requirements of Swine, National Research Council. Nutrient requirements of swine. 11th ed. Washington, DC:

- National Academy Press; 2012.
2. Cromwell GL, Stahly TS, Monegue HJ. Effects of source and level of copper on performance and liver copper stores in weanling pigs. *J Anim Sci* 1989;67:2996-3002.
 3. Mei SF, Yu B, Ju CF, Zhu D, Chen DW. Effect of different levels of copper on growth performance and cecal ecosystem of newly weaned piglets. *Ital J Anim Sci* 2010;9:378-81.
 4. Coffey RD, Cromwell GL, Monegue HJ. Efficacy of a copper-lysine complex as a growth promotant for weanling pigs. *J Anim Sci* 1994;72:2880-6.
 5. Zhao J, Allee G, Gerlemann G, et al. Effects of a chelated copper as growth promoter on performance and carcass traits in pigs. *Asian-Australas J Anim Sci* 2014;27:965-73.
 6. Hellman NE, Gitlin JD. Ceruloplasmin metabolism and function. *Annu Rev Nutr* 2002;22:439-58.
 7. Dove C. Copper citrate as a growth stimulating copper source for nursery pigs. *J Anim Sci* 1998;76:159.
 8. Armstrong TA, Cook DR, Ward MM, Williams CM, Spears JW. Effect of dietary copper source (cupric citrate and cupric sulfate) and concentration on growth performance and fecal copper excretion in weanling pigs. *J Anim Sci* 2004;82:1234-40.
 9. Bremner I. Manifestations of copper excess. *Am J Clin Nutr* 1998;67:1069S-73S.
 10. Harris ZL, Takahashi Y, Miyajima H, et al. Aceruloplasminemia: molecular characterization of this disorder of iron-metabolism. *Proc Natl Acad Sci USA* 1995;92:2539-43.
 11. Amer M, Elliot J. Effects of supplemental dietary copper on glyceride distribution in the backfat of pigs. *Can J Anim Sci* 1973;53:147-52.
 12. Amer M, Elliot J. Effects of level of copper supplement and removal of supplemental copper from the diet on the physical and chemical characteristics of porcine depot fat. *Can J Anim Sci* 1973;53:139-45.
 13. Massie HR, Ofosuappiah W, Aiello VR. Elevated serum copper is associated with reduced immune response in aging mice. *Gerontology* 1993;39:136-45.
 14. Percival SS. Copper and immunity. *Am J Clin Nutr* 1998;67:1064S-8S.
 15. Hill G, Miller E, Whetter P, Ullrey D. Concentration of minerals in tissues of pigs from dams fed different levels of dietary zinc. *J Anim Sci* 1983;57:130-8.
 16. Huang YL, Ashwell MS, Fry RS, et al. Effect of dietary copper amount and source on copper metabolism and oxidative stress of weanling pigs in short-term feeding. *J Anim Sci* 2015;93:2948-55.
 17. Li J, Yan L, Zheng X, et al. Effect of high dietary copper on weight gain and neuropeptide Y level in the hypothalamus of pigs. *J Trace Elem Med Biol* 2008;22:33-8.
 18. Zhu D, Yu B, Ju C, Mei S, Chen D. Effect of high dietary copper on the expression of hypothalamic appetite regulators in weanling pigs. *J Anim Feed Sci* 2011;20:60-70.
 19. Ma YL, Zanton GI, Zhao J, et al. Multitrial analysis of the effects of copper level and source on performance in nursery pigs. *J Anim Sci* 2015;93:606-14.
 20. Armstrong TA, Spears JW, van Heugten E, Engle TE, Wright CL. Effect of copper source (cupric citrate vs cupric sulfate) and level on growth performance and copper metabolism in pigs. *Asian-Australas J Anim Sci* 2000;13:1154-61.
 21. Kim BE, Nevitt T, Thiele DJ. Mechanisms for copper acquisition, distribution and regulation. *Nat Chem Biol* 2008;4:176-85.
 22. Pastorelli G, Rossi R, Zanardi E, Ghidini S, Corino C. Two different forms and levels of CuSO₄ in piglet feeding: liver, plasma and faeces copper status. *J Anim Feed Sci* 2014;23:52-7.
 23. Cater MA, La Fontaine S, Shield K, Deal Y, Mercer JFB. ATP7B mediates vesicular sequestration of copper: Insight into biliary copper excretion. *Gastroenterology* 2006;130:493-506.
 24. Kaim W, Rall J. Copper-a "modern" bioelement. *Angew Chem Int Edit* 1996;35:43-60.
 25. Ward JD, Spears JW. Long-term effects of consumption of low-copper diets with or without supplemental molybdenum on copper status, performance, and carcass characteristics of cattle. *J Anim Sci* 1997;75:3057-65.
 26. Senthilkumar P, Nagalakshmi D, Reddy YR, Sudhakar K. Effect of different level and source of copper supplementation on immune response and copper dependent enzyme activity in lambs. *Trop Anim Health Prod* 2009;41:645-53.
 27. Draper H, Hadley M. Malondialdehyde determination as index of lipid peroxidation. *Methods Enzymol* 1989;186:421-31.
 28. Fry RS, Ashwell MS, Lloyd KE, et al. Amount and source of dietary copper affects small intestine morphology, duodenal lipid peroxidation, hepatic oxidative stress, and mRNA expression of hepatic copper regulatory proteins in weanling pigs. *J Anim Sci* 2012;90:3112-9.
 29. Feng J, Ma WQ, Gu ZL, Wang YZ, Liu JX. Effects of dietary copper (II) sulfate and copper proteinate on performance and blood indexes of copper status in growing pigs. *Biol Trace Elem Res* 2007;120:171-8.
 30. Armstrong TA, Spears JW, Engle TE, See MT. Effect of pharmacological concentrations of dietary copper on lipid and cholesterol metabolism in pigs. *Nutr Res* 2001;21:1299-308.
 31. Li QH, Luo X, Liu B, Han J, Yu S. The effect of dietary copper as copper-glycine on blood physiological and biochemical traits and tissue copper contents of weanling pigs. *Acta Vet Zootech Sin* 2004;35:23-7 (In chinese with english abstract).
 32. Bonham M, O'Connor JM, Hannigan BM, Strain J. The immune system as a physiological indicator of marginal copper status? *Br J Nutr* 2002;87:393-403.
 33. Kelley DS, Daudu PA, Taylor PC, Mackey BE, Turnlund JR. Effects of low-copper diets on human immune response. *Am J Clin Nutr* 1995;62:412-6.
 34. Wang XX, Song PX, Wu H, et al. Effects of graded levels of isomaltoligosaccharides on the performance, immune function and intestinal status of weaned pigs. *Asian-Australas J Anim Sci* 2016;29:250-6.
 35. Yan JY, Zhang C, Tang L, Kuang SY. Effect of dietary copper sources and concentrations on serum lysozyme concentration and protegrin-1 gene expression in weaning piglets. *Ital J Anim Sci* 2015;14:3709.

36. Espinoza A, Le Blanc S, Olivares M, et al. Iron, copper, and zinc transport: inhibition of divalent metal transporter 1 (DMT1) and human copper transporter 1 (hCTR1) by shRNA. *Biol Trace Elem Res* 2012;146:281-6.
37. Collins JF, Prohaska JR, Knutson MD. Metabolic crossroads of iron and copper. *Nutr Rev* 2010;68:133-47.
38. Vasak M, Meloni G. Chemistry and biology of mammalian metallothioneins. *J Biol Inorg Chem* 2011;16:1067-78.
39. Pang Y, Applegate TJ. Effects of dietary copper supplementation and copper source on digesta pH, calcium, zinc, and copper complex size in the gastrointestinal tract of the broiler chicken. *Poult Sci* 2007;86:531-7.